





**EPIPEN Auto-Injector and EPIPEN Jr Auto-Injector Directions**

- **First, remove the EPIPEN Auto-Injector from the plastic carrying case**
- **Pull off the blue safety release cap**



- **Hold orange tip near outer thigh (always apply to thigh)**



- **Swing and firmly push orange tip against outer thigh. Hold on thigh for approximately 10 seconds. Remove the EPIPEN Auto-Injector and massage the area for 10 more seconds**



DEY® and the Dey logo, EpiPen®, EpiPen 2-Pak®, and EpiPen Jr 2-Pak® are registered trademarks of Dey Pharma, L.P.

**An allergy response kit should contain at least two doses of epinephrine, other medications as noted by the student’s physician, and a copy of this Allergy Action Plan. A kit must accompany the student if he/she is off school grounds (i.e., field trip).**

**CONTACTS: 911**

**LOCAL EMERGENCY CONTACTS:**

**NAME/RELATIONSHIP**

**PHONE (\_\_\_\_) \_\_\_\_\_**

**PHONE (\_\_\_\_) \_\_\_\_\_**

**NAME/RELATIONSHIP**

**PHONE (\_\_\_\_) \_\_\_\_\_**

**PHONE (\_\_\_\_) \_\_\_\_\_**